

# Sum-Enchanted Evenings

The Fun and Joy of Mathematics



## LECTURE 6

**Peter Lynch**

**School of Mathematics & Statistics  
University College Dublin**

**Evening Course, UCD, Autumn 2018**



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability

Astronomy I

Numbers

The Number Line



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability

Astronomy I

Numbers

The Number Line



# Meaning and Content of Mathematics

The word **Mathematics** comes from Greek *μαθημα* (*máthéma*), meaning “knowledge” or “study” or “learning”.

It is the study of topics such as

- ▶ Quantity (numbers)
- ▶ Structure (patterns)
- ▶ Space (geometry)
- ▶ Change (analysis).



# Outline

Introduction

**Axioms and Proof**

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability

Astronomy I

Numbers

The Number Line



# What are Axioms?

How can we prove a theorem,  
if we have nothing to start from?

We cannot prove something using nothing.  
We need some starting point.

The basic building blocks are called **Axioms**.

Axioms are not proved, but are assumed true.



# What are Axioms?

**Axioms are important because the entire body of mathematics rests upon them.**

If there are **too few axioms**, we can prove very little of interest from them.

If there are **too many axioms**, we can prove almost any result from them.

**Consistency:**

**We must not have axioms that contradict each other.**



# What are Axioms?

**Mathematicians assume that axioms are true without being able to prove them.**

**This is not problematic, because axioms are normally intuitively obvious.**





# What are Axioms?

Mathematicians assume that axioms are true without being able to prove them.

This is not problematic, because axioms are normally intuitively obvious.

There are usually only a few axioms. For example, we may assume that

$$a \times b = b \times a$$

for any two numbers  $a$  and  $b$ .



# What are Axioms?

Mathematicians assume that axioms are true without being able to prove them.

This is not problematic, because axioms are normally intuitively obvious.

There are usually only a few axioms. For example, we may assume that

$$a \times b = b \times a$$

for any two numbers  $a$  and  $b$ .

But Hamilton found that for **quaternions**,

$$A \times B \neq B \times A.$$



**Different sets of axioms lead to different kinds of mathematics.**

**Every area of mathematics has its own set of basic axioms.**



**Different sets of axioms lead to different kinds of mathematics.**

**Every area of mathematics has its own set of basic axioms.**

**When mathematicians have proven a theorem, they publish it for other mathematicians to check.**

**In principle, it is possible to break a proof into steps starting from the basic axioms.**



**Different sets of axioms lead to different kinds of mathematics.**

**Every area of mathematics has its own set of basic axioms.**

**When mathematicians have proven a theorem, they publish it for other mathematicians to check.**

**In principle, it is possible to break a proof into steps starting from the basic axioms.**

**Sometimes a mistake in the proof is found.**

**Sometimes an error is not found for many years (e.g., an early “proof” of the **Four Colour Theorem.**)**



# Euclid's Axioms of Geometry

Euclid based his “Elements of Geometry” on a set of five postulates or axioms:

"Let the following be postulated":

1. "To draw a **straight line** from any **point** to any point."
2. "To produce [extend] a **finite straight line** continuously in a straight line."
3. "To describe a **circle** with any centre and distance [radius]."
4. "That all **right angles** are equal to one another."
5. *The parallel postulate*: "That, if a straight line falling on two straight lines make the interior angles on the same side less than two right angles, the two straight lines, if produced indefinitely, meet on that side on which are the angles less than the two right angles."

The fifth postulate, the **parallel postulate**, has been a great source of controversy and confusion. This has led to **completely new areas of mathematics**.



# Peano's Axioms of Arithmetic

Giuseppe Peano constructed five axioms to build up the set  $\mathbb{N}$  of natural numbers:

$$\exists 0 : 0 \in \mathbb{N}$$

$$\forall n \in \mathbb{N} : \exists n' \in \mathbb{N}$$

$$\neg(\exists n \in \mathbb{N} : n' = 0)$$

$$\forall m, n \in \mathbb{N} : m' = n' \Rightarrow m = n$$

$$\forall A \subseteq \mathbb{N} : (0 \in A \wedge (n \in A \Rightarrow n' \in A)) \Rightarrow A = \mathbb{N}$$

The natural numbers may then be extended to the integers, rational numbers and real numbers.



# Axioms of Set Theory

Set theory is the basic language of mathematics.

Many mathematical problems can be formulated in the language of set theory.

To prove them we need the **Set Theory Axioms**.


The most widely accepted axioms are the set of **nine Zermelo-Fraenkel (ZF) axioms**.

A tenth axiom, may also be assumed, the **Axiom of Choice**.





# Zermelo-Fraenkel axioms



## AXIOM OF EXTENSION

If two sets have the same elements, then they are equal.




## AXIOM OF SEPERATION

We can form a subset of a set, which consists of some elements.




## EMPTY SET AXIOM

There is a set with no members, written as  $\{\}$  or  $\emptyset$ .




## PAIR-SET AXIOM

Given two objects  $x$  and  $y$  we can form a set  $\{x, y\}$ .



## UNION AXIOM

We can form the union of two or more sets.



## POWER SET AXIOM

Given any set, we can form the set of all subsets (the power set).

Image from Mathigon.org



# Zermelo-Fraenkel axioms



## AXIOM OF INFINITY

There is a set with infinitely many elements.




## AXIOM OF FOUNDATION

Sets are built up from simpler sets, meaning that every (non-empty) set has a minimal member.



## AXIOM OF REPLACEMENT

If we apply a function to every element in a set, the answer is still a set.



## AXIOM OF CHOICE

Given infinitely many non-empty sets, you can choose one element from each of these sets.

Image from Mathigon.org



# Axiom of Choice




Image from Wikipedia



# Axiom of Choice

The Axiom of Choice (AC) looks just as innocuous as the other nine axioms.  
**However it has unexpected consequences.**

We can use AC to prove that it is possible to cut a sphere into five pieces and reassemble them into two spheres, each identical to the initial sphere.



# Axiom of Choice

The Axiom of Choice (AC) looks just as innocuous as the other nine axioms. **However it has unexpected consequences.**

We can use AC to prove that it is possible to cut a sphere into five pieces and reassemble them into two spheres, each identical to the initial sphere.

This result is called the **Banach-Tarski Theorem**.



# Banach-Tarski Theorem



The five pieces have fractal boundaries:  
they can't actually be made in practice.

Also, they are not **measurable**:  
they have no definite volume.



# The Current Status

**There is an active debate among logicians about whether to accept the Axiom of Choice or not.**

**Every collection of axioms forms a different “mathematical world”. Different theorems may be true in different worlds.**

**The question is: are we happy to live in a world where we can make two spheres from one.**

[See Wikipedia article: Axiom of Choice](#)



# Outline

Introduction

Axioms and Proof

**Three Utilities Problem**

Greek 4

Distraction 12: Conditional Probability

Astronomy I

Numbers


The Number Line






# Three Utilities Problem: Abstract

Is the complete  $3 \times 3$  bipartite graph  $K_{3,3}$  planar?



# Three Utilities Problem: Abstract

Is the complete  $3 \times 3$  bipartite graph  $K_{3,3}$  planar?



This is an abstract, jargon-filled question in **topological graph theory**.


We look at a simple, concrete version.



# Three Utilities Problem: Concrete

We have to connect 3 utilities to 3 houses.


- ▶ Electricity
- ▶ Water
- ▶ Gas




The lines must not cross.



# Three Utilities Problem: Have a Go




# Three Utilities Problem: Solution!



[http://www.archimedes-lab.org/How\\_to\\_Solve/Water\\_gas.html](http://www.archimedes-lab.org/How_to_Solve/Water_gas.html)




# Three Utilities Problem: No Solution!




[http://www.archimedes-lab.org/How\\_to\\_Solve/Water\\_gas.html](http://www.archimedes-lab.org/How_to_Solve/Water_gas.html)



# Three Utilities Problem



# Three Utilities Problem: Application






# Three Utilities Problem for Mugs



# Three Utilities Problem on a Torus





$K_{3,3}$  is a toroidal graph.

Vi Hart: <https://www.youtube.com/watch?v=CruQy1WSfoU&feature=youtu.be&t=9>





# Three Utilities: Kuratowski's Theorem

If a graph contains  $K_{3,3}$  or  $K_5$  as a sub-graph, it is **non-planar**. If it does not contain either, it is **planar**.



# Three Utilities: Equivalent Graphs



The two forms shown are equivalent.

There are crossings in both.



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

**Greek 4**

Distraction 12: Conditional Probability

Astronomy I

Numbers

The Number Line



# The Greek Alphabet, Part 4

α	β	γ	δ	ε	ζ
Alpha	Beta	Gamma	Delta	Epsilon	Zeta
η	θ	ι	κ	λ	μ
Eta	Theta	Iota	Kappa	Lambda	Mu
ν	ξ	ο	π	ρ	σ
Nu	Xi	Omicron	Pi	Rho	Sigma
τ	υ	φ	χ	ψ	ω
Tau	Upsilon	Phi	Chi	Psi	Omega

Figure : 24 beautiful letters



# The Last Six Letters

We will consider the final group of six letters.

$\tau$        $\upsilon$        $\phi$        $\chi$        $\psi$        $\omega$

T      Υ      Φ      Χ      Ψ      Ω

Let us focus first on the **small letters**  
and come back to the big ones later.



$\tau$     $\upsilon$     $\phi$     $\chi$     $\psi$     $\omega$

**Tau: You have certainly heard of a Tau-cross:  $\tau$ .**

**Upsilon ( $\upsilon$ ) or u-psilon means ‘bare u’.  
It is often transliterated as ‘y’.**

**Phi ( $\phi$ ) is ‘f’, often used for latitude  
(as  $\lambda$  is often used for longitude).**

**Chi ( $\chi$ ) has a ‘ch’ or ‘k’ sound.**

**Psi ( $\psi$ ) is very common: psychology, etc.**

**Omega ( $\omega$ ) is the end: Alpha and Omega  $\left(\frac{\text{A}}{\Omega}\right)$ .**





$\tau$     $\upsilon$     $\phi$     $\chi$     $\psi$     $\omega$

**Tau: You have certainly heard of a Tau-cross:  $\tau$ .**

**Upsilon ( $\upsilon$ ) or u-psilon means ‘bare u’.  
It is often transliterated as ‘y’.**

**Phi ( $\phi$ ) is ‘f’, often used for latitude  
(as  $\lambda$  is often used for longitude).**

**Chi ( $\chi$ ) has a ‘ch’ or ‘k’ sound.**

**Psi ( $\psi$ ) is very common: psychology, etc.**

**Omega ( $\omega$ ) is the end: Alpha and Omega  $\left(\frac{\text{A}}{\Omega}\right)$ .**

**Now you know 24 letters. You should get a diploma.**



# A Few Greek Words (for practice)

κωμα

ψυκη

κρισις

αναθεμα

αμβροσια

καταστροφη



# A Few Greek Words (for practice)

κωμα

ψυκη

κρισις

αναθεμα

αμβροσια

καταστροφη

**Coma:** κωμα

**Psyche:** ψυκη

**Crisis:** κρισις

**Anathema:** αναθεμα

**Ambrosia:** αμβροσια

**Catastrophe:** καταστροφη









# End of Greek 104



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

**Distraction 12: Conditional Probability**

Astronomy I

Numbers

The Number Line






# Distraction 12: Conditional Probability

## Conditional Probability

Conditional Probability: Level 3 Challenges



A box contains two white marbles and two black marbles. I pick a marble at random and set it aside. Then, I pick a second marble and notice that it is black.

Is it more likely that the first marble was white or black?



# Distraction 12: Conditional Probability

Possible outcomes of the experiment:

$$W_1 W_2 \quad W_1 B_2 \quad B_1 W_2 \quad B_1 B_2$$

Are all four possibilities equally likely?



# Distraction 12: Conditional Probability

**Possible outcomes of the experiment:**

$$W_1 W_2 \quad W_1 B_2 \quad B_1 W_2 \quad B_1 B_2$$

**Are all four possibilities equally likely?**

$$P(B_2) = P(W_1)P(B_2|W_1) + P(B_1)P(B_2|B_1)$$

$$P(W_1) = \frac{1}{2} \quad P(B_1) = \frac{1}{2} \quad P(B_2|W_1) = \frac{2}{3} \quad P(B_2|B_1) = \frac{1}{3}$$



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability


**Astronomy I**

Numbers

The Number Line



# The Quadrivium



## The Pythagorean model of mathematics



# The Ancient Greeks

**Mathematics and Astronomy are intimately linked.**

**Two of the strands of the Quadrivium were  
Geometry (static) and Cosmology (dynamic space).**

**Greek astronomer Claudius Ptolemy (c.90–168AD)  
placed the Earth at the centre of the universe.**

**The Sun and planets move around the Earth in orbits  
that are of the most perfect of all shapes: circles.**



# Aristarchus of Samos (c.310–230 BC)


Aristarchus of Samos (*Ἀριστάρχος*), astronomer and mathematician, presented the first model that placed the Sun at the center of the universe.

The original writing of Aristarchus is lost, but Archimedes wrote in his **Sand Reckoner**:

*“His hypotheses are that the fixed stars and the Sun remain unmoved, that the Earth revolves about the Sun on the circumference of a circle, ... ”*




# Eratosthenes (c.276–194 BC)






# Eratosthenes (c.276–194 BC)



# Eratosthenes (c.276–194 BC)



# Hipparchus (c.190–120 BC)

**Hipparchus of Nicaea (*Ἰππάρχος*) was a Greek astronomer, geographer, and mathematician.**

**Regarded as the greatest astronomer of antiquity.**

**Often considered to be the founder of trigonometry.**

**Famous for**

- ▶ **Precession of the equinoxes**
- ▶ **First comprehensive star catalog**
- ▶ **Invention of the astrolabe**
- ▶ **Invention (perhaps) of the armillary sphere.**



# Claudius Ptolemy (c.AD 100–170)

Claudius Ptolemy was a Greco-Roman astronomer, mathematician, geographer and astrologer.

He lived in the city of Alexandria.

Ptolemy wrote several scientific treatises:


- ▶ An astronomical treatise (the *Almagest*) originally called *Mathematical Treatise (Mathematike Syntaxis)*.
- ▶ A book on geography.
- ▶ An astrological treatise.

Ptolemy's *Almagest* is the only surviving comprehensive ancient treatise on astronomy.



# Ptolemy's Model

Ptolemy's model was universally accepted until the appearance of simpler heliocentric models during the scientific revolution.



# Epicycles Rule

According to **Norwood Russell Hanson**  
(science historian):


*There is no bilaterally symmetrical, nor eccentrically periodic curve used in any branch of astrophysics or observational astronomy which could not be smoothly plotted as the resultant motion of a point turning within a constellation of epicycles, finite in number, revolving around a fixed deferent.*

“The Mathematical Power of Epicyclical Astronomy”, 1960


**Any path — periodic or not, closed or open — can be represented by an infinite number of epicycles.**



# Ptolemaic Epicycles



# Conic Sections



Circles are special cases  
of **conic sections**.

They are formed by a plane  
cutting a cone at an angle.

Conics were studied by **Apollonius of Perga**  
(late 3rd – early 2nd centuries BC).

[https://en.wikipedia.org/wiki/Conic\\_section](https://en.wikipedia.org/wiki/Conic_section)






# The Scientific Revolution

## TRAILER

Next week, we will look at developments in the sixteenth and seventeenth centuries.



Nicolaus Copernicus  
1473 – 1543



Tycho Brahe  
1546 – 1601



Johannes Kepler  
1571 – 1630



Galileo Galilei  
1564 – 1642



Figure from [mathigon.org](http://mathigon.org)



# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability

Astronomy I

**Numbers**

The Number Line



# Babylonian Numerals

𐎶 1	𐎠𐎺 11	𐎠𐎶𐎶 21	𐎠𐎶𐎶𐎶 31	𐎠𐎶𐎶𐎶𐎶 41	𐎠𐎶𐎶𐎶𐎶𐎶 51
𐎶𐎶 2	𐎠𐎶𐎶 12	𐎠𐎶𐎶𐎶 22	𐎠𐎶𐎶𐎶𐎶 32	𐎠𐎶𐎶𐎶𐎶𐎶 42	𐎠𐎶𐎶𐎶𐎶𐎶𐎶 52
𐎶𐎶𐎶 3	𐎠𐎶𐎶𐎶 13	𐎠𐎶𐎶𐎶𐎶 23	𐎠𐎶𐎶𐎶𐎶𐎶 33	𐎠𐎶𐎶𐎶𐎶𐎶𐎶 43	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶 53
𐎶𐎶𐎶𐎶 4	𐎠𐎶𐎶𐎶𐎶 14	𐎠𐎶𐎶𐎶𐎶𐎶 24	𐎠𐎶𐎶𐎶𐎶𐎶𐎶 34	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶 44	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 54
𐎶𐎶𐎶𐎶𐎶 5	𐎠𐎶𐎶𐎶𐎶𐎶 15	𐎠𐎶𐎶𐎶𐎶𐎶𐎶 25	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶 35	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 45	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 55
𐎶𐎶𐎶𐎶𐎶𐎶 6	𐎠𐎶𐎶𐎶𐎶𐎶𐎶 16	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶 26	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 36	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 46	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 56
𐎶𐎶𐎶𐎶𐎶𐎶𐎶 7	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶 17	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 27	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 37	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 47	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 57
𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 8	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 18	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 28	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 38	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 48	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 58
𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 9	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 19	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 29	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 39	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 49	𐎠𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶𐎶 59
𐎶𐎶𐎶 10	𐎠𐎶𐎶 20	𐎠𐎶𐎶𐎶 30	𐎠𐎶𐎶𐎶𐎶 40	𐎠𐎶𐎶𐎶𐎶𐎶 50	



# Ancient Egyptian Numerals

1 =		10 =	∩	100 =	☉	1000 =	𐎗
2 =		20 =	∩∩	200 =	☉☉	2000 =	𐎗𐎗
3 =		30 =	∩∩∩	300 =	☉☉☉	3000 =	𐎗𐎗𐎗
4 =		40 =	∩∩∩∩	400 =	☉☉☉☉	4000 =	𐎗𐎗𐎗𐎗
5 =		50 =	∩∩∩∩∩	500 =	☉☉☉☉☉	5000 =	𐎗𐎗𐎗𐎗𐎗



# Ancient Hebrew and Greek Numerals





















8 𐤇 Chet ח	7 ז Zayin ז	6 ו Vav ו	5 ה Hey ה	4 ד Dalet ד	3 ג Gimmel ג	2 ב Bet ב	1 א Aleph א
70 ע Ayin ע	60 ס Samekh ס	50 נ Nun נ	40 מ Mem מ	30 ל Lamed ל	20 כ Kaf כ	10 י Yod י	9 ט Tet ט

1	$\alpha$	alpha	10	$\iota$	iota	100	$\rho$	rho
2	$\beta$	beta	20	$\kappa$	kappa	200	$\sigma$	sigma
3	$\gamma$	gamma	30	$\lambda$	lambda	300	$\tau$	tau
4	$\delta$	delta	40	$\mu$	mu	400	$\upsilon$	upsilon
5	$\epsilon$	epsilon	50	$\nu$	nu	500	$\phi$	phi
6	$\zeta$	vau*	60	$\xi$	xi	600	$\chi$	chi
7	$\zeta$	zeta	70	$\omicron$	omicron	700	$\psi$	psi
8	$\eta$	eta	80	$\pi$	pi	800	$\omega$	omega
9	$\theta$	theta	90	$\koppa^*$	koppa*	900	$\sampi$	sampi

\*vau, koppa, and sampi are obsolete characters



# Mayan Numerals

 0	 1	 2	 3	 4
 5	 6	 7	 8	 9
 10	 11	 12	 13	 14
 15	 16	 17	 18	 19



# Various Numeral Systems

## Numeral systems

**0123456789**

·|١٢٣٤٥٦٧٨٩

I II III IV V VI VII VIII IX X

○ १ २ ३ ४ ५ ६ ७ ८ ९

○ ൧ ൨ ൩ ൪ ൫ ൬ ൭ ൮ ൯

○ ୦ ୧ ୨ ୩ ୪ ୫ ୬ ୭ ୮ ୯

**○ 一 二 三 四 五 六 七 八 九**

Wikipedia: Hindu-Arabic Numeral System



# Roman Numerals

I	1	XXI	21	XLI	41
II	2	XXII	22	XLII	42
III	3	XXIII	23	XLIII	43
IV	4	XXIV	24	XLIV	44
V	5	XXV	25	XLV	45
VI	6	XXVI	26	XLVI	46
VII	7	XXVII	27	XLVII	47
VIII	8	XXVIII	28	XLVIII	48
IX	9	XXIX	29	XLIX	49
X	10	XXX	30	L	50
XI	11	XXXI	31	LI	51
XII	12	XXXII	32	LII	52
XIII	13	XXXIII	33	LIII	53
XIV	14	XXXIV	34	LIV	54
XV	15	XXXV	35	LV	55
XVI	16	XXXVI	36	LVI	56
XVII	17	XXXVII	37	LVII	57
XVIII	18	XXXVIII	38	LVIII	58
XIX	19	XXXIX	39	LIX	59
XX	20	XL	40	LX	60

In order:  $MDC LXVI = 1666$





# How to Multiply Roman Numbers

**Table :** Multiplication Table for Roman Numbers.

	<b>I</b>	<b>V</b>	<b>X</b>	<b>L</b>	<b>C</b>	<b>D</b>	<b>M</b>
<b>I</b>	<i>I</i>	<i>V</i>	<i>X</i>	<i>L</i>	<i>C</i>	<i>D</i>	<i>M</i>
<b>V</b>	<i>V</i>	<i>XXV</i>	<i>L</i>	<i>CCL</i>	<i>D</i>	<i>MMD</i>	$\overline{V}$
<b>X</b>	<i>X</i>	<i>L</i>	<i>C</i>	<i>D</i>	<i>M</i>	$\overline{V}$	$\overline{X}$
<b>L</b>	<i>L</i>	<i>CCL</i>	<i>D</i>	<i>MMD</i>	$\overline{V}$	$\overline{XXV}$	$\overline{L}$
<b>C</b>	<i>C</i>	<i>D</i>	<i>M</i>	$\overline{V}$	$\overline{X}$	$\overline{L}$	$\overline{C}$
<b>D</b>	<i>D</i>	<i>MMD</i>	$\overline{V}$	$\overline{XXV}$	$\overline{L}$	$\overline{CCL}$	$\overline{D}$
<b>M</b>	<i>M</i>	$\overline{V}$	$\overline{X}$	$\overline{L}$	$\overline{C}$	$\overline{D}$	$\overline{M}$



# A Roman Abacus


Replica of a Roman abacus from 1st century AD.




Abacus is a Latin word, which comes from the Greek  $\alpha\beta\alpha\kappa\alpha\varsigma$  (board or table).




# A Chinese Abacus: Suan Pan



# A Japanese Abacus: Soroban




# A Different Angle on Numerals



**Delightful theory. Almost certainly wrong.**






## Arguments “for”

1. It is a very simple idea
2. It links symbols to numerical values






## Arguments “for”

1. It is a very simple idea
2. It links symbols to numerical values

## Arguments “against”

1. Number forms modified to fit model
2. Complete lack of historical evidence.





## Arguments “for”

1. It is a very simple idea
2. It links symbols to numerical values

## Arguments “against”

1. Number forms modified to fit model
2. Complete lack of historical evidence.

The great tragedy of science —

the slaying of a beautiful hypothesis by an ugly fact (T H Huxley)





# Outline

Introduction

Axioms and Proof

Three Utilities Problem

Greek 4

Distraction 12: Conditional Probability

Astronomy I

Numbers

**The Number Line**



# A Hierarchy of Numbers

We will introduce a hierarchy of numbers.

Each set is contained in the next one.

They are like a set of nested Russian Dolls:



Matryoshka

# The Natural Numbers $\mathbb{N}$

The **counting numbers** were the first to emerge:

1 2 3 4 5 6 7 8 ...

They are also called the **Natural Numbers**.



# The Natural Numbers $\mathbb{N}$

The **counting numbers** were the first to emerge:

1 2 3 4 5 6 7 8 ...

They are also called the **Natural Numbers**.



We can arrange the natural numbers in a list.

This list is like a **toy computer**.



# The Natural Numbers $\mathbb{N}$

The set of natural numbers is denoted  $\mathbb{N}$ .

If  $n$  is a natural number, we write  $n \in \mathbb{N}$ .



# The Natural Numbers $\mathbb{N}$

The set of natural numbers is denoted  $\mathbb{N}$ .

If  $n$  is a natural number, we write  $n \in \mathbb{N}$ .

Natural numbers can be added:  $4 + 2 = 6 \in \mathbb{N}$



But not always subtracted:  $4 - 6 = -2 \notin \mathbb{N}$ .



# The Natural Numbers $\mathbb{N}$

The set of natural numbers is denoted  $\mathbb{N}$ .

If  $n$  is a natural number, we write  $n \in \mathbb{N}$ .

Natural numbers can be added:  $4 + 2 = 6 \in \mathbb{N}$



But not always subtracted:  $4 - 6 = -2 \notin \mathbb{N}$ .

To allow for subtraction **we have to extend**  $\mathbb{N}$ .



# The Integers $\mathbb{Z}$

We extend the counting numbers by adding the negative whole numbers:

... -3 -2 -1 0 1 2 3 4 ...

The whole numbers are also called the **Integers**.





# The Integers $\mathbb{Z}$

We extend the counting numbers by adding the negative whole numbers:

... -3 -2 -1 0 1 2 3 4 ...

The whole numbers are also called the **Integers**.

The set of integers is denoted  $\mathbb{Z}$ .

If  $k$  is an integer, we write  $k \in \mathbb{Z}$ .

Clearly,

$$\mathbb{N} \subset \mathbb{Z}$$



**Integers can be added and subtracted.**

**They can also multiplied:**

$$6 \times 4 = 24 \in \mathbb{Z}.$$



**Integers can be added and subtracted.**

**They can also multiplied:**

$$6 \times 4 = 24 \in \mathbb{Z}.$$

**However, they cannot usually be divided:**

$$\frac{6}{4} = 1\frac{1}{2} \notin \mathbb{Z}.$$



**Integers can be added and subtracted.**

**They can also multiplied:**

$$6 \times 4 = 24 \in \mathbb{Z}.$$

**However, they cannot usually be divided:**

$$\frac{6}{4} = 1\frac{1}{2} \notin \mathbb{Z}.$$

**To allow for division we have to extend  $\mathbb{Z}$ .**



# The Rational Numbers $\mathbb{Q}$

We extend the integers by adding fractions:

$$r = \frac{p}{q} \quad \text{where } p \text{ and } q \text{ are integers.}$$

These **rational numbers** are **ratios of integers**.



# The Rational Numbers $\mathbb{Q}$

We extend the integers by adding fractions:

$$r = \frac{p}{q} \quad \text{where } p \text{ and } q \text{ are integers.}$$

These **rational numbers** are **ratios of integers**.

The set of rational numbers is denoted  $\mathbb{Q}$ .

If  $r$  is a rational number, we write  $r \in \mathbb{Q}$ .

Clearly,

$$\mathbb{Z} \subset \mathbb{Q}$$



With the Rational Numbers, we can:

**Add, Subtract, Multiply and Divide**

That is, for any  $p \in \mathbb{Q}$  and  $q \in \mathbb{Q}$

All of  $p + q$   $p - q$   $p \times q$  and  $p \div q$

are rational numbers.



With the Rational Numbers, we can:

**Add, Subtract, Multiply and Divide**

That is, for any  $p \in \mathbb{Q}$  and  $q \in \mathbb{Q}$

All of  $p + q$   $p - q$   $p \times q$  and  $p \div q$

are rational numbers.

We say that  $\mathbb{Q}$  is **closed under addition, subtraction, multiplication and division.**





With the Rational Numbers, we can:

**Add, Subtract, Multiply and Divide**

That is, for any  $p \in \mathbb{Q}$  and  $q \in \mathbb{Q}$

All of  $p + q$   $p - q$   $p \times q$  and  $p \div q$


are rational numbers.

We say that  $\mathbb{Q}$  is **closed under addition, subtraction, multiplication and division.**

But we are not yet finished.  $\mathbb{R}$  is yet to come.



# The Hierarchy of Numbers



$N \subset Z \subset Q \subset R \subset C$



# The Hierarchy of Numbers

Each set is contained in the next one.

They are like a set of nested Russian Dolls:



Matryoshka

Thank you

